

# Dietary Fat Is Associated With Metabolic Syndrome in Japanese Brazilians

RENATA D. FREIRE<sup>1</sup>  
MARLY A. CARDOSO, PHD<sup>2</sup>  
SUELY G.A. GIMENO, PHD<sup>3</sup>

SANDRA R.G. FERREIRA, MD, PHD<sup>3</sup>  
FOR THE JAPANESE-BRAZILIAN DIABETES  
STUDY GROUP\*

**OBJECTIVE** — The aim of this study was to examine the association of nutrient intakes with metabolic syndrome in a Japanese descendant population at high risk for metabolic abnormalities.

**RESEARCH DESIGN AND METHODS** — In a population-based study of Japanese Brazilians aged  $\geq 30$  years, 412 men and 465 women were studied. The diagnosis of metabolic syndrome was based on the National Cholesterol Education Program criteria modified for Asians. Food intake was assessed by a validated food frequency questionnaire.

**RESULTS** — Men and women showed similar mean ages ( $55.7 \pm 12.7$  and  $54.4 \pm 11.9$  years) and 49% (95% CI 44.8–54.6) and 43% (38.4–47.6) had metabolic syndrome, respectively. As expected, clinical parameters were less favorable in the subset of subjects with metabolic syndrome. Men with metabolic syndrome showed higher cholesterol ( $233.2 \pm 116.3$  vs.  $211.7 \pm 5.8$  g/day,  $P < 0.05$ ) and lower carbohydrate ( $288.5 \pm 45.8$  vs.  $300.1 \pm 39.8$  g/day,  $P < 0.001$ ) intakes than those without metabolic syndrome, but no difference was observed among the women. After adjusting for sex, age, smoking, education level, generation, physical activity, total energy, and dietary fiber intake, a positive association between metabolic syndrome and total fat intake was detected. Comparing people in the highest quintile of total fat consumption with those in the lowest quintile, odds ratio (OR) of metabolic syndrome was 5.0 [95% CI 1.58–16.00];  $P < 0.005$ . In contrast, linoleic acid intake was inversely associated with metabolic syndrome (OR 0.50 [95% CI 0.26–0.98];  $P < 0.05$ ). Considering food groups, after adjustments only fried food intake was shown to be associated with increased risk of metabolic syndrome.

**CONCLUSIONS** — Despite the limitation of such a study design in investigating cause-effect relationships, our findings favor the hypothesis that dietary total fat may increase whereas linoleic acid intake may reduce the risk of metabolic syndrome in Japanese descendants living in Brazil.

*Diabetes Care* 28:1779–1785, 2005

The concept of metabolic syndrome includes a number of metabolic disturbances linked by insulin resistance, which increase cardiovascular risk (1). Weight gain (in particular, the abdominal depot of body fat) has a key role for the development of the syndrome.

Lack of physical activity and certain dietary patterns, including high saturated fatty acid and low vegetable intake, contribute to weight gain and increase the risk of metabolic disturbances (2,3), whereas such potentially modifiable lifestyle factors may reduce cardiovascular risk.

From the <sup>1</sup>Department of Internal Medicine, Division of Endocrinology, Federal University of São Paulo, São Paulo, Brazil; the <sup>2</sup>Department of Nutrition, School of Public Health, University of São Paulo, São Paulo, Brazil; and the <sup>3</sup>Department of Preventive Medicine, Federal University of São Paulo, São Paulo, Brazil.

Address correspondence and reprint requests to Dr. Sandra R.G. Ferreira, Universidade Federal de São Paulo, Dept. Medicina Preventiva, Rua Botucatu, 740, São Paulo, SP, Brazil 04023-062. E-mail: ferreira@medprev.epm.br.

Received for publication 28 February 2005 and accepted in revised form 5 April 2005.

\*A complete list of Japanese-Brazilian Diabetes Study Group members can be found in the APPENDIX.

**Abbreviations:** HOMA-IR, homeostasis model assessment of insulin resistance.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

© 2005 by the American Diabetes Association.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Metabolic syndrome is highly prevalent in the U.S. with an estimated 24% of the adult population affected (4). The National Cholesterol Education Program (5) defined metabolic syndrome based on several parameters including anthropometric measurements. However, anthropometric cutoffs to identify high-risk subjects may be influenced by ethnicity (6,7). Based on the fact that in Asian populations morbidity and mortality are occurring in people with lower BMI values and waist circumferences, the World Health Organization recently proposed different cutoff values to define increased morbidity in Asians (7).

Migration studies indicate that adoption of the Western lifestyle is strongly associated with type 2 diabetes. Our group conducted a population-based study in Japanese migrants and their descendants in Brazil who have one of the highest prevalence of glucose metabolism disturbances worldwide (8). High prevalence rates of type 2 diabetes and associated diseases in Japanese migrants living in the Americas (8–11) suggest that the exposure to a different lifestyle exacerbated an inherent tendency to accumulate fat and develop diabetes (12). Our group and others reported that dietary changes, such as an increase in animal fat intake and reduction in complex carbohydrates, were imposed by immigration (12–15).

Animal studies suggested that the type of fat in diet may affect insulin sensitivity by changing the fatty acid composition of membrane lipids (16). In humans, a detrimental effect of saturated fat or a benefit of polyunsaturated fat is controversial (3,17–19). Few data are available regarding the association of dietary factors with clusters of diseases that have a common nutritional basis, such as the metabolic syndrome (20). Considering the complexity of establishing relationships between diet and disease, the focus of this study was on a population in which several diseases cluster together. Therefore, we examined the association of nutrient intakes with metabolic syndrome in a Japanese-Brazilian population at high risk for metabolic abnormalities.

## RESEARCH DESIGN AND METHODS

A survey was designed to estimate the prevalence and incidence of diabetes and associated diseases in a Japanese-Brazilian population living in Bauru, São Paulo, Brazil (8). This was approved by the Institutional Ethics Committee and written informed consent was obtained from all participants. The entire Japanese-Brazilian population  $\geq 30$  years of age ( $n = 1,651$ ) was invited, and 1,330 first-generation (Japan-born) and second-generation (Brazil-born) subjects participated in all phases of the survey. Self-reported diabetes, hypertension, dyslipidemia, and other diseases whose treatment could interfere in the dietary plan were excluded. Thus, results refer to 877 eligible subjects. They were interviewed at home by trained interviewers using standardized questionnaires and scheduled for physical examination and laboratory procedures after overnight fasting. The protocol has been described in detail elsewhere (8).

### Dietary data

Dietary data were obtained by a quantitative food frequency questionnaire, developed and validated for the Japanese-Brazilian population (21). Subjects were asked about their usual consumption of foods and food groups (122 items) during the last year. Participants reported their habitual frequency of consumption of each item, the respective unit of time (if daily, weekly, monthly, or annually), and size of the usual individual portion (if small, average, large, or extra-large in relation to the reference portion for each item of the food frequency questionnaire). Questions concerning use of sauce, frequency of intake of visible fat, and type of fat used in cooking procedures were also included. Complete rechecking of the coding as well as double keying were performed on every questionnaire. In addition to standard range edits, internal consistency edits and nutrient calculations for the diets were performed using Dietsys 4.01 software (22). The nutrient database used was based primarily on U.S. Department of Agriculture publications supplied by Dietsys, supplemented by the most recent edition of Standard Food Composition Tables of Brazil (23) and Japan (24). The variables selected for this study were total calories, carbohydrate, protein, total fat, saturated fatty acid, oleic acid, linoleic acid, chole-

sterol, total fiber, and alcohol. Macronutrients were also considered as percentages of total calories.

Physical activity was assessed by a questionnaire including items regarding the types of physical activities performed during work and leisure periods. A score was attributed to each subject, and three categories were created: light, moderate, or heavy. Smoking habit was classified into two categories: current smoker or not. Education level was classified into three categories ( $<1$ , 1–8, and  $>8$  years) and generation as Japan-born or Brazil-born subjects. Body weight and height were measured while subjects wore light clothing and no shoes using calibrated electronic scales and a fixed stadiometer. Waist circumference was measured at the umbilicus. BMI was calculated as weight (kilograms) divided by height (meters) squared. Blood pressure was taken three times by automatic device (Omron model HEM-712C; Omron Health Care) and the mean value of the two last measurements was used to express systolic and diastolic blood pressure.

### Laboratory methods

Total cholesterol, its fractions, triglyceride, and uric acid were measured enzymatically by automatic analyzer. Samples for insulin determination were stored at  $-80^{\circ}\text{C}$  until assay; this was determined by monoclonal antibody-based immunofluorimetric assay (AutoDelfia; Perkin Elmer Life Sciences, Norton, OH). Homeostasis model assessment of insulin resistance (HOMA-IR) was calculated according to Matthews et al. (25).

Diagnosis of metabolic syndrome was based on the Third Report of the National Cholesterol Education Program—Adult Treatment Panel III (5), replacing the anthropometric cutoffs by those previously proposed by the World Health Organization for Asians (7). The presence of three or more of the following components defined metabolic syndrome: fasting plasma glucose  $\geq 110$  mg/dl (6.1 mmol/l); systolic or diastolic blood pressure  $\geq 130$  or  $\geq 85$  mmHg; HDL cholesterol  $< 50$  mg/dl (1.29 mmol/l) for women or  $< 40$  mg/dl (1.04 mmol/l) for men; triglyceride  $\geq 150$  mg/dl (1.79 mmol/l); and waist circumference  $\geq 80$  for women or  $\geq 90$  cm for men.

### Statistical methods

Statistical analysis was performed using SPSS version 10.0 (SPSS, Chicago, IL). Rates of metabolic syndrome were calculated by point and interval. Nutrient intake was expressed in grams and as percentages of total energy adjusted for total energy through the residual method. Energy-adjusted nutrient intakes were calculated as the residuals from the regression model, with absolute nutrient intake as the dependent variable and total energy intake as the independent variable (26). Nutritional, clinical, and laboratory data of the participants with and without metabolic syndrome were compared by Student's *t* test; for categorical variables, the Mantel-Haenszel  $\chi^2$  test for trend was applied. Logistic regression was used to calculate the odds ratios (ORs) and their 95% CIs for metabolic syndrome, with individuals in the lowest quintile category of macronutrients as the reference category. ORs were adjusted for sex, age, smoking, generation, leisure physical activity, level of education, total energy, dietary fiber, and alcohol intake and for a number of possible sex interactions. To assess trends across quintile categories, we assigned the median intake of each quintile category to individuals with intakes in the category and then included this quintile median variable as a continuous factor in logistic regression models. The *P* for trend was the resulting *P* value for the associated logistic regression coefficient (27). Statistical significance was defined as a two-tailed *P* value  $< 0.05$ .

**RESULTS**— Similar proportions of men (47%,  $n = 412$ ) and women (53%,  $n = 465$ ) were included, and their mean ages were  $55.7 \pm 12.7$  and  $54.4 \pm 11.9$  years, respectively. Slightly higher prevalence of metabolic syndrome was found in men (49.8% [95% CI 44.8–54.6]) than in women (43.0% [38.4–47.6]). As expected, clinical and biochemical parameters were less favorable, and HOMA-IR values were higher in the subsets of Japanese-Brazilian men and women with metabolic syndrome (Table 1). Proportions of generations, current smokers, and physical activity levels did not differ between subjects with or without metabolic syndrome.

Nutrient intakes according to sex and the presence of metabolic syndrome are shown in Table 2. Men with metabolic syndrome had significantly higher total

Table 1—Main characteristics of the study participants

	Men		Women	
	Without metabolic syndrome	With metabolic syndrome	Without metabolic syndrome	With metabolic syndrome
<i>n</i>	207	205	265	200
Age (years)	55.2 ± 13.7	56.3 ± 11.7	52.4 ± 12.2	57.2 ± 10.8†
Generation (%)				
Japan born	51 (24.6)	37 (18.0)	29 (10.9)	28 (14.0)
Brazil born	156 (75.4)	168 (82.0)	236 (89.1)	172 (86.0)
Education level (%)				
<1 year	25 (12.1)	16 (7.8)	23 (8.7)	27 (13.5)†
1–8 years	76 (36.7)	84 (41.0)	103 (39.0)	112 (56.0)
>8 years	106 (51.2)	105 (51.2)	138 (52.3)	61 (30.5)
Current cigarette smoking (%)				
Yes	44 (21.3)	34 (16.7)	24 (9.1)	20 (10.0)
No	163 (78.7)	170 (83.3)	241 (90.9)	180 (90.0)
Physical activity (%)				
Light	162 (78.3)	161 (78.5)	223 (84.5)	172 (86.0)
Moderate	39 (18.8)	40 (19.5)	36 (13.6)	25 (12.5)
Heavy	6 (2.9)	4 (2.0)	5 (1.9)	3 (1.5)
BMI (kg/m <sup>2</sup> )	23.1 ± 2.8	26.3 ± 3.4†	22.4 ± 2.8	25.8 ± 3.3†
Waist (cm)	82.1 ± 7.2	91.1 ± 8.1†	74.0 ± 6.8	83.1 ± 7.9†
Systolic blood pressure (mmHg)	122.4 ± 18.3	139.5 ± 20.1†	115.5 ± 17.1	135.3 ± 22.8†
Diastolic blood pressure (mmHg)	73.6 ± 10.0	84.4 ± 11.7†	70.2 ± 9.3	79.0 ± 12.1†
Plasma glucose (mmol/l)	6.24 ± 0.75	7.17 ± 1.48†	5.96 ± 0.62	6.88 ± 1.76†
Insulin (pmol/l)	36.6 ± 24.0	63.0 ± 45.6†	40.8 ± 31.2	57.0 ± 34.8†
HOMA-IR	1.7 ± 1.8	3.3 ± 2.4†	1.8 ± 1.5	2.9 ± 2.0†
Total cholesterol (mmol/l)	5.32 ± 0.96	5.53 ± 1.02*	5.35 ± 1.06	5.74 ± 1.05†
LDL cholesterol (mmol/l)	3.29 ± 0.94	3.33 ± 0.95	3.28 ± 0.9	3.52 ± 0.97*
HDL cholesterol (mmol/l)	1.29 ± 0.27	1.23 ± 0.30*	1.38 ± 0.27	1.29 ± 0.26*
Triglycerides (mmol/l)	2.26 ± 1.84	3.72 ± 2.92†	1.68 ± 1.33	3.04 ± 2.25†
Uric acid (mmol/l)	0.38 ± 0.09	0.43 ± 0.10†	0.27 ± 0.06	0.32 ± 0.07†

Data are means ± SD. \* $P < 0.01$ , † $P < 0.001$  vs. without metabolic syndrome.

fat (in grams and percentage of total energy), oleic acid, and cholesterol intakes than those without metabolic syndrome ( $P < 0.05$ ). In women, significantly lower protein intake adjusted for total energy was observed when comparing subsets with and without the syndrome.

ORs of metabolic syndrome across quintiles of dietary nutrients showed that the highest quintile of total fat was associated with a fivefold increase in risk of having the syndrome, after adjustment for sex, age, smoking, education level, generation, physical activity, total energy intake, dietary fiber, alcohol, and sex times years of education interaction (Table 3). Conversely, the highest consumption of linoleic acid was associated with a protection against metabolic syndrome (OR 0.50 [95% CI 0.26–0.98]).

Nutrients were also considered in food groups such as quintiles of fruits, bread/cereals/rice, red meat, fish, eggs, and oils/fried food. After adjustment for

sex, age, smoking, education, generation, physical activity, total energy, total fat, and linoleic acid, only fried food intake was shown to be associated with increased risk of metabolic syndrome (OR 1.27 [95% CI 0.82–1.96], 1.42 [0.92–2.19], 1.31 [0.85–2.04], and 1.69 [1.10–2.61] for the second, third, fourth, and fifth quintiles, respectively;  $P = 0.02$ ).

**CONCLUSIONS**— Although aspects of diet have been linked to individual metabolic features of the metabolic syndrome (2,3,17,18), the role of dietary factors in the etiopathogenesis of this syndrome has been poorly investigated (20). As far as type 2 diabetes is concerned, conflicting relationships with carbohydrates and fat intake have been reported (28,29). Grouping diseases that share a common etiopathogenetic basis was an attempt to clarify the relationships of the insulin resistance syndrome with dietary factors. We had an opportunity to study a

population of Japanese ancestry that was shown to be highly susceptible to metabolic syndrome when exposed to the Western world (30). Since 1993, our group has followed a cohort of Japanese Brazilians with high prevalence of central obesity, diabetes, dyslipidemia, and hypertension (8,9,12).

The definition of metabolic syndrome used in the present study introduced a modification to the National Cholesterol Education Program criteria with respect to the waist circumference cutoffs (5), in accordance with recent World Health Organization recommendations for Asian populations (7). The same diagnostic approach was previously used in other Asian populations, namely the Omani from Saudi Arabia (31). A slightly higher proportion of Japanese-Brazilian men with metabolic syndrome are compatible with greater cardiovascular risk attributed to sex. Consistently higher HOMA-IR values in the subsets of Japanese-Brazilian men

Table 2—Daily intake of energy and selected nutrients of the study participants

	Men		Women	
	Without metabolic syndrome	With metabolic syndrome	Without metabolic syndrome	With metabolic syndrome
<i>n</i>	207	205	265	200
Energy (kcal)	2177 ± 615	2293 ± 670	1868 ± 518	1823 ± 542
Carbohydrate				
Crude intake (g)	293.0 ± 84.4	295.6 ± 95.4	248.2 ± 75.4	246.2 ± 83.2
Total energy adjusted (g)	300.1 ± 39.8	288.5 ± 45.8*	245.6 ± 32.4	249.7 ± 32.2
Total energy (%)	54.2 ± 7.5	51.7 ± 7.8†	53.1 ± 7.1	53.8 ± 7.1
Protein				
Crude intake (g)	72.0 ± 24.5	76.5 ± 25.3	65.3 ± 23.5	61.4 ± 20.5
Total energy adjusted (g)	73.9 ± 13.4	74.6 ± 14.5	64.7 ± 13.7	62.3 ± 11.5†
Total energy (%)	13.2 ± 2.4	13.3 ± 2.5	13.9 ± 2.7	13.5 ± 2.4
Total fat				
Crude intake (g)	76.5 ± 29.9	83.7 ± 29.5†	69.9 ± 22.6	67.7 ± 22.7
Total energy adjusted (g)	78.7 ± 15.6	81.4 ± 16.6	69.2 ± 12.0	68.6 ± 12.7
Total energy (%)	31.3 ± 6.2	32.7 ± 6.1†	33.7 ± 6.1	33.5 ± 6.4
Oleic acid (g)				
Crude intake (g)	28.5 ± 12.0	31.4 ± 11.3†	26.5 ± 8.7	25.4 ± 8.7
Total energy adjusted (g)	29.3 ± 7.4	30.6 ± 7.2	26.3 ± 5.6	25.8 ± 5.8
Total energy (%)	11.7 ± 2.8	12.3 ± 2.8†	12.9 ± 2.8	12.6 ± 2.9
Linoleic acid (g)				
Crude intake (g)	12.0 ± 5.6	12.8 ± 5.0	11.4 ± 4.0	11.0 ± 4.3
Total energy adjusted (g)	12.4 ± 3.6	12.4 ± 3.2	11.3 ± 2.9	11.2 ± 3.2
Total energy (%)	4.9 ± 1.4	5.0 ± 1.2	5.5 ± 1.4	5.5 ± 1.5
Saturated fatty acid (g)				
Crude intake (g)	18.5 ± 7.8	20.4 ± 8.3†	16.8 ± 6.5	16.3 ± 6.6
Total energy adjusted (g)	19.1 ± 4.6	19.8 ± 5.2	16.6 ± 4.1	16.6 ± 4.3
Total energy (%)	7.5 ± 1.8	7.9 ± 2.0	8.0 ± 2.0	8.0 ± 2.1
Cholesterol (mg)				
Crude intake (g)	205.0 ± 95.6	239.4 ± 139.8†	175.1 ± 87.5	167.3 ± 81.0
Total energy adjusted (g)	211.1 ± 78.8	233.2 ± 116.3†	173.2 ± 70.4	169.8 ± 63.1
Fiber (g)				
Crude intake (g)	18.2 ± 7.7	18.9 ± 9.0	16.0 ± 6.5	16.3 ± 7.0
Total energy adjusted (g)	18.7 ± 5.8	18.4 ± 6.7	15.9 ± 4.9	16.5 ± 5.4
Alcohol‡ (g)				
Crude intake (g)	138.2 ± 266.3	193.6 ± 341.1	9.6 ± 31.2	12.3 ± 53.4
Total energy adjusted (g)	143.4 ± 264.7	188.5 ± 333.4	9.6 ± 31.2	12.3 ± 53.3

Data are means ± SD. \* $P < 0.001$ , † $P < 0.05$  vs. without metabolic syndrome. ‡Log-transformed values for statistical purposes.

and women classified with metabolic syndrome assured a reasonable identification of subjects with metabolic syndrome in whom associations with nutrient intake were investigated.

Although food frequency techniques have been largely used in cross-sectional surveys, they could be considered limited by their lack of quantitative accuracy. Underreporting of energy intake may be a problem when obese subjects are under investigation. The term “underreporting” applies to dietary assessment methods used to estimate total energy intake, such as the food frequency questionnaire. The small number of Japanese Brazilians with elevated BMI should minimize the

possibility of systematic bias caused by underreporting in the present study. Additionally, as pointed out by Margetts and Nelson (32), differences in mean intake levels of macronutrients between underreporters and those who give valid records are reduced by energy adjustment.

In crude analysis, men with metabolic syndrome were shown to have higher total fat and cholesterol intake to the detriment of carbohydrate than those without the syndrome. However, no impact on the risk of metabolic syndrome was shown across quintiles of carbohydrate intake adjusted for a number of variables. The relationship between dietary carbohydrates and insulin sensitivity is a matter of

controversy (28,33). Simple sugars may have a detrimental effect (33), whereas starches and polysaccharides may have benefits for glucose metabolism (20,34,35). In a large sample of adult Swedish men and women, food patterns dominated by fiber bread provided favorable effects, whereas those high in refined bread or in cheese, cake, and alcoholic beverages increased the risk for several components of metabolic syndrome (36). Concordantly, recent data from the Framingham Offspring Study supported the fact that whole-grain intake is inversely associated with HOMA-IR and lower prevalence of metabolic syndrome (20). Our study and others provided conflict-



Table 3—ORs of metabolic syndrome across quintiles of energy-adjusted nutrient intake

	Quintiles of intake					P value
	1 (lowest)	2	3	4	5 (highest)	
Protein						
Median intake (g/day)	51.9	56.2	60.9	70.2	91.3	
Range of intake (g/day)	17.4–150.3	22.2–107.6	30.4–119.3	28.4–133.5	44.0–185.6	
OR	1.00	0.57 (0.34–0.94)	0.72 (0.42–1.22)	0.62 (0.35–1.09)	0.69 (0.38–1.27)	0.448
Total fat						
Median intake (g/day)	52.9	61.4	67.5	76.9	95.6	
Range of intake (g/day)	16.2–175.8	17.5–129.1	25.5–139.3	40.3–145.1	47.1–249.8	
OR	1.00	0.63 (0.32–1.22)	1.39 (0.62–3.08)	2.15 (0.81–5.69)	5.03 (1.58–16.0)	0.005
Oleic acid						
Median intake (g/day)	20.0	22.5	24.5	29.7	36.7	
Range of intake (g/day)	6.6–52.6	6.0–42.2	9.4–51.7	14.7–74.3	20.0–113.7	
OR	1.00	0.85 (0.44–1.64)	0.89 (0.41–1.91)	0.71 (0.29–1.70)	0.72 (0.25–2.06)	0.426
Linoleic acid						
Median intake (g/day)	7.9	9.4	10.1	11.9	15.4	
Range of intake (g/day)	2.43–17.8	3.1–19.4	3.8–20.8	3.7–23.1	7.5–45.7	
OR	1.00	0.92 (0.52–1.62)	1.08 (0.65–1.81)	0.65 (0.34–1.21)	0.50 (0.26–0.98)	0.026
Saturated fatty acid						
Median intake (g/day)	12.2	13.8	15.3	17.6	24.3	
Range of intake (g/day)	3.7–45.6	5.4–39.8	4.3–31.7	8.8–35.5	11.9–62.7	
OR	1.00	0.87 (0.52–1.46)	1.18 (0.67–2.06)	1.01 (0.55–1.85)	0.77 (0.39–1.50)	0.480
Cholesterol (mg)						
Median intake (mg/day)	109.6	136.6	163.1	206.4	297.2	
Range of intake (mg/day)	16.9–298.0	45.2–509.1	35.4–300.3	84.9–353.2	140.3–931.3	
OR	1.00	1.40 (0.86–2.29)	1.40 (0.79–2.48)	1.13 (0.61–2.07)	1.36 (0.72–2.55)	0.900

Data are OR (95% CI) unless indicated and are adjusted for sex, age ( $\leq 45$ , 46–60, and  $> 60$  years), smoking, years of education ( $< 1$ , 1–8, and  $> 8$  years), generation, physical activity, total energy intake, dietary fiber, alcohol, and interaction (sex times years of education).

ing results which could be attributed to the heterogeneity of study designs and/or source of dietary carbohydrates.

The lack of association of carbohydrate intake seen in the present study is somewhat in agreement with the study by Brunner et al. (37). However, these investigators did not detect considerable differences between types of dietary fat, which was the case in our study. Despite this, OR was seen as a measurement of risk for metabolic syndrome in the present study. Because OR may overestimate the risk of a common outcome such as metabolic syndrome, it is possible that an increase in risk attributed to fat intake by the OR is, in fact, less pronounced than a fivefold increase. Our findings of an adverse impact of total fat intake and protection of polyunsaturated fat concerning the association with metabolic syndrome are supported by several investigations (2,38,39). Some human studies have specifically investigated the relationship of linoleic acid intake and metabolic syndrome-related disorders and reinforced a protection role for such polyunsaturated acid (40–44). In both muscle and adipose tissue, a high-fat diet resulted in de-

creased insulin receptor numbers, but no change in receptor affinity. The same diet has been shown to decrease insulin-stimulated glucose transport and intracellular glucose metabolism (45). The type of fat and degree of saturation were also shown to play an important role for glucose metabolism. Classically, saturated fatty acids and cholesterol have been positively associated with fasting and postprandial glucose levels (46). Several studies indicated the fatty acid composition of membranes is dependent on the fatty acid composition of diet (47). Diets high in polyunsaturated fatty acid result in increased insulin receptor numbers but decreased receptor affinity. On the other hand, saturated fat could exert adverse effects on the glucose/insulin system by mechanisms involving oxidative stress.

Among the environmental changes imposed on the Japanese immigrants in Brazil, changes in dietary habits were probably one of the most important. Their current dietary pattern contrasts with the traditional Japanese diet, particularly concerning the proportion of fat, which, in association with sedentary lifestyle, may be contributing to increased

visceral adiposity and a predisposition to metabolic syndrome.

In conclusion, our data suggest that total fat intake may be a risk factor for metabolic syndrome in a Japanese-Brazilian population. Our findings also corroborate a beneficial effect of polyunsaturated fatty acid such as linoleic acid to reduce risk for this clustering of diseases linked by insulin resistance. Prospective studies will help test the hypothesis in this population.

**Acknowledgments**— This study was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo.

## APPENDIX

### Members of Japanese-Brazilian Diabetes Study Group

Alcides Hirai, Amélia T. Hirai, Helena Harima, Magid Iunes (in memoriam), Mário Kikuchi, Katsumi Osiro, Sandra R. G. Ferreira, and Suely G. A. Gimeno (Department of Preventive Medicine, Federal University of São Paulo, Brazil); Laércio J. Franco (Department of Social Medicine,

Faculty of Medicine of Ribeirão Preto, University of São Paulo, Brazil); Regina C. S. Moisés and Luiza Matsumura (Department of Internal Medicine, Federal University of São Paulo, Brazil); Newton de Barros Jr. (Department of Surgery, Federal University of São Paulo, Brazil); Vania D'Almeida (Pediatrics Department, Federal University of São Paulo, Brazil); Nilce Tomita (Dental School of Bauru, University of São Paulo, Brazil); Katsunori Wakisaka (Japanese-Brazilians Study Center, Brazil); Rita Chaim (Department of Nutrition, University of Sagrado Coração de Jesus, Bauru, Brazil); and Marly Augusto Cardoso (Department of Nutrition, School of Public Health, University of São Paulo, Brazil).

# References

1. Lakka HM, Laaksonen DE, Lakka TA, Niiskanen LK, Kumpusalo E, Tuomilehto J, Salonen JT: The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. *JAMA* 288:2709–2716, 2002
2. van Dam RM, Willett WC, Rimm EB, Stampfer MJ, Hu FB: Dietary fat and meat intake in relation to risk of type 2 diabetes in men. *Diabetes Care* 25:417–424, 2002
3. Feskens EJ, Virtanen SM, Rasanen L, Tuomilehto J, Stengard J, Pekkanen J, Nissinen A, Kromhout D: Dietary factors determining diabetes and impaired glucose tolerance: a 20-year follow-up of the Finnish and Dutch cohorts of the Seven Countries Study. *Diabetes Care* 18:1104–1112, 1995
4. Ford ES, Giles WH, Dietz WH: Prevalence of the metabolic syndrome among US adults. *JAMA* 287:356–359, 2002
5. Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP). Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA* 285:2486–2497, 2001
6. Kasim-Karakas SE: Ethnic differences in the insulin resistance syndrome. *Am J Clin Nutr* 71:670–671, 2000
7. Inoue S, Zimmet P, Bassett J: *The Asia-Pacific Perspective: Redefining Obesity and its Treatment*. Melbourne, Australia, World Health Organization Health, 2000
8. Gimeno SGA, Ferreira SRG, Cardoso MA, Franco LJ, Hirai AT, Matsumura L, Moisés RS for the Japanese-Brazilians Diabetes Study Group. Prevalence and 7-year incidence of type 2 diabetes mellitus in a Japanese-Brazilian population: an alarming public health problem. *Diabetologia* 45: 1635–1638, 2002
9. Ferreira SRG, Iunes M, Franco LJ, Iochida LC, Hirai A, Vivolo MA for the Japanese-Brazilians Diabetes Study Group. Disturbances of glucose and lipid metabolism in first and second generation of Japanese-Brazilians. *Diabetes Res Clin Pract* 34:S59–S63, 1996
10. Fujimoto WY, Leonetti DL, Kinyoun JL, Newell-Morris L, Shuman WP, Storov WC, Wahl PW: Prevalence of diabetes mellitus and impaired glucose tolerance among second generation Japanese-American men. *Diabetes* 36:721–729, 1987
11. Hara H, Egusa G, Yamakido M, Kawate R: The high prevalence of diabetes mellitus and hyperinsulinemia among the Japanese-Americans living in Hawaii and Los Angeles. *Diabetes Res Clin Pract* 24:S37–S42, 1994
12. Ferreira SRG, Lerario DDG, Gimeno SGA, Sanudo A, Franco LJ: Obesity and central adiposity in Japanese immigrants: role of the Western dietary pattern. *J Epidemiol* 12:431–438, 2002
13. Costa MB, Ferreira SRS, Franco LJ, Gimeno SGA, Iunes M: Dietary patterns in a high-risk population for glucose intolerance. *J Epidemiol* 10:111–117, 2000
14. Freire RD, Cardoso MA, Shinzato AR, Ferreira SRG: Nutritional status of Japanese-Brazilian subjects: comparison across gender and generation. *Br J Nutr* 89:705–712, 2003
15. Tsunehara CH, Leonetti DL, Fujimoto WY: Diet of second-generation Japanese-American men with and without non-insulin-dependent diabetes. *Am J Clin Nutr* 52:731–738, 1990
16. Storlien LH, Baur LA, Kriketos AD, Pan Clooney GJ, Jenkins AB, Calvert GD, Campbell LV: Dietary fats and insulin action. *Diabetologia* 39:621–631, 1996
17. Marshall JA, Bessesen DH, Hamman RF: High saturated fat and low starch and fibre are associated with hyperinsulinaemia in a non-diabetic population: the San Luis Valley Diabetes Study. *Diabetologia* 40: 430–438, 1997
18. Trevisan M, Krogh V, Freudenheim J, Blake A, Muti P, Panico S, Farinero E, Mancini M, Menotti A, Ricci G: Consumption of olive oil, butter, and vegetable oils and coronary heart disease risk factors: the Research Group ATS-RF2 of the Italian National Research Council. *JAMA* 263:688–692, 1990
19. Mayer-Davis EJ, Monaco JH, Hoen HM, Carmichael S, Vitolins MZ, Rewers MJ, Haffner SM, Ayad MF, Bergman RN, Karter AJ: Dietary fat and insulin sensitivity in a triethnic population: the role of obesity: the Insulin Resistance Atherosclerosis Study (IRAS). *Am J Clin Nutr* 65: 79–87, 1997
20. McKeon NM, Meigs JB, Liu S, Saltzman E, Wilson PWF, Jacques PF: Carbohydrate nutrition, insulin resistance, and the prevalence of the metabolic syndrome in the Framingham Offspring Cohort. *Diabetes Care* 27:538–546, 2004
21. Cardoso MA, Kida AA, Tomita LY, Stocco PR: Reproducibility and validity of a food frequency questionnaire among women of Japanese ancestry living in Brazil. *Nutr Res* 21:725–733, 2001
22. Block G, Coyle LM, Hartman AM, Scoppa metabolic syndrome: Revision of dietary analysis software for the health habits and history questionnaire. *Am J Epidemiol* 139: 1190–1196, 1994
23. Fundação IBGE. *Tabela de composição de alimentos*. 4th ed. Rio de Janeiro, IBGE, 1996
24. Resources Council, Science and Technology Agency. *Standard Tables of Food Composition in Japan*. 4th ed. Tokyo, Women's University of Nutrition Press, 1993
25. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC: Homeostasis model assessment: insulin resistance and  $\beta$ -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28:412–419, 1985
26. Willett WC, Stampfer MJ: Total energy intake: implications for epidemiological analyses. *Am J Epidemiol* 124:17–27, 1986
27. Kleinbaum DG, Kupper LL, Muller KE: *Applied Regression Analysis and Other Multivariate Methods*. Boston, PWS-Kent Publishing Company, 1988
28. Feskens EJM: Nutritional factors and the etiology of non-insulin-dependent diabetes mellitus: an epidemiological overview. *World Rev Nutr Diet* 96:1–39, 1992
29. Mooy JM, Grootenhuys PA, de Vries H, Valkenburg HA, Bouter LM, Kostense PJ, Heine RJ: Prevalence and determinants of glucose intolerance in a Dutch Caucasian population: the Hoorn Study. *Diabetes Care* 18:1270–1273, 1995
30. Rosenbaum P, Gimeno SGA, Sanudo A, Franco LJ, Ferreira SRG: Analysis of criteria for metabolic syndrome in a population-based study of Japanese-Brazilians. *Diabetes Obes Metab*. In press
31. Al-Lawati JA, Mohammed AJ, Al-Hinai HQ, Jousilahti P: Prevalence of the metabolic syndrome among Omani adults. *Diabetes Care* 26:1781–1785, 2003
32. Margetts BM, Nelson M: *Desing Concepts in Nutritional Epidemiology*. Oxford, Oxford University Press, 1998
33. Daly ME, Vale CV, Walker M, Alberti KG, Mathers: Dietary carbohydrates and insulin sensitivity: a review of evidence and clinical implications. *Am J Clin Nutr* 66: 1072–1085, 1997
34. Salmeron J, Manson JE, Stampfer MJ, Colditz GA, Wing AL, Willett WC: Dietary fiber, glycemic load, and risk of non-insulin-dependent diabetes mellitus in

- women. *J Am Med Assoc* 277:472–477, 1997
35. Meyer KA, Kushi LH, Jacobs DR, Slavin J, Sellers TA, Folsom AR: Carbohydrates dietary fiber, and incident type 2 diabetes in older women. *Am J Clin Nutr* 71:921–930, 2000
  36. Wirfält E, Hedblad Bo, Gullberg Bo, Mattisson I, Andrén C, Rosander U, Janzon L, Berglund G: Food patterns and components of the metabolic syndrome in men and women: a cross-sectional study within the Malmö Diet and Cancer Cohort. *Am J Epidemiol* 154:1150–1159, 2001
  37. Brunner EJ, Wunsch H, Marmot MG: What is an optimal diet? Relationship of macronutrient intake to obesity, glucose tolerance, lipoprotein cholesterol levels and the metabolic syndrome in the Whitehall II study. *Int J Obes* 25:45–53, 2001
  38. Feskens EJM, Bowles CH, Kromhout D: Inverse association between fish intake and risk of glucose intolerance in normoglycemic elderly men and women. *Diabetes Care* 14:935–941, 1991
  39. Salmeron J, Hu FB, Manson JE, Stampfer MJ, Colditz GA, Rimm EB, Willett WC: Dietary fat intake and risk of type 2 diabetes in women. *Am J Clin Nutr* 73:1019–1026, 2001
  40. Laaksonen DE, Lakka TA, Lakka HM, Nyyssönen K, Rissanen T, Niskanen LK, Salonen JT: Serum fatty acid composition predicts development of impaired fasting glycaemia and diabetes in middle-aged men. *Diabet Med* 19:456–464, 2002
  41. Djoussé L, Folsom AR, Province MA, Hunt SC, Ellison RC: Dietary linoleic acid and carotid atherosclerosis: the National Heart, Lung, and Blood Institute Family Heart Study. *Am J Clin Nutr* 77:819–825, 2003
  42. Djoussé L, Arnett DK, Pankow JS, Hopkins PN, Province MA, Ellison RC: Dietary linolenic acid is associated with a lower prevalence of hypertension in the NHLBI Family Heart Study. *Hypertension* 45:368–373, 2005
  43. Iso H, Sato S, Umemura U, Kudo M, Koibe K, Kitamura A, Imano H, Okamura T, Naito Y, Shimamoto T: Linoleic acid, other fatty acids, and the risk of stroke. *Stroke* 33:2086–2093, 2002
  44. Wang L, Folsom AR, Zheng ZJ, Pankow JS, Eckfeldt JH: Plasma fatty acid composition and incidence of diabetes in middle-aged adults: the Atherosclerosis Risk in Communities (ARIC) Study. *Am J Clin Nutr* 78:91–98, 2003
  45. Fields M: Nutritional factors adversely influencing the glucose/insulin system. *Am J Clin Nutr* 17:317–321, 1998
  46. Beck-Nielsen H, Pederson O, Schwartz SN: Effects of diet on the cellular insulin binding and the insulin sensitivity in young healthy subjects. *Diabetologia* 15: 289–299, 1978
  47. Tepperman HM, Dewitt J, Tepperman J: Hormone effects on glycogenesis, glucogenesis, and cyclic AMP production by liver cells from rats fed diets high in glucose or lard. *J Nutr* 108:1924–1928, 1978